


Office: 555 Prince Charles Drive N

Welland, ON L3C 4J6 Phone: 905-808-3103 Fax: 905-228-2955

**Email:** info@drjedrzejko.com **Web:** www.drjedrzejko.com

# Hemorrhoid Banding/ Injection Instructions

After Your Endoscopic Procedure



#### **Preparation**

Purchase a rectal Fleet enema from your pharmacy. This must be used 2hr prior to your procedure.

#### What is expected in the first 48-72hr?

- Slight bleeding with or without bowel movements
- Some discomfort in the bum/perineum
- Sensation of needing to have a bowel movement
- Some bleeding after 7-12 days when the hemorrhoid sloughs off

## What should you do?

- Avoid constipation
- Simple analgesia (e.g., Tylenol, Advil)
- Hot water sitz baths
- Banding may need to be repeated in 4-6 weeks if symptoms persist

## When should you see a doctor?

- Excessive pain
- Excessive bleeding per rectum
- Difficulty passing urine
- Fevers/chills

## **Managing pain**

It is normal to have some discomfort or pain. This should slowly improve but might last a few days to weeks. Your pain should still allow you to move, deep breathe, poop, and do every-day tasks.

Contact Dr. Jedrzejko if you would like to book an in-person or phone follow-up appointment 2-3 weeks after your procedure.

